

Precautions After Filler

1. Immediately after the procedure, bruising and swelling will vary by individual, and the treated area may appear uneven.
2. Swelling usually lasts for about 3–4 days, and you may feel a burning or heated sensation in the treated area.
3. Be careful of external pressure, as it can change the shape of the treated area. (For lip or corner-of-mouth procedures, avoid frequent straw use and ensure hot foods do not directly touch the area.)
4. For 1 month after treatment, avoid alcohol, smoking, saunas, and intense exercise that causes facial heat.
5. Avoid rubbing or applying strong stimulation to the treated area.
6. Stitches (if present) can be removed after 7 days.
7. After stitch removal, showering is allowed. Until then, wash your face with a damp towel, avoiding the stitched area. (If there are no stitches, you may shower starting from the evening of the following day.)
8. After the procedure, common side effects may include swelling, redness, mild pain, increased sensitivity, or discoloration at the injection site.
 - These symptoms usually subside within 2 days.
 - If symptoms persist, worsen, or if the area becomes increasingly painful, swollen, or dark in color, seek medical attention immediately.

⚠ If you experience any unusual discomfort or have further questions, please contact the clinic.



www.okayps.co



+82 10 3395 6887



contact@okayps.co



4F, 5F, 715 Samseong-ro, Gangnam-gu, Seoul



WhatsApp